

ETDRS LOGMAR SEHTAFELTEST

PATIENT:

RE

LI

Rx: SPH _____ CYL _____ Achse _____ <input type="checkbox"/> Brille <input type="checkbox"/> Kontaktlinse <input type="checkbox"/> IOL <input type="checkbox"/> zentrale Sehschärfe	Anzahl Fehler	Deci	Snellen 20ft	LogMar @ 4 M	Snellen Equivalent	VA %	Anzahl Fehler	Rx: SPH _____ CYL _____ Achse _____ <input type="checkbox"/> Brille <input type="checkbox"/> Kontaktlinse <input type="checkbox"/> IOL <input type="checkbox"/> zentrale Sehschärfe
NCKZO		0.10	20/200	1.0	6/60	20%		DSRKN
RHSDK		0.12	20/160	0.9	6/48	28%		CKZOH
DOVHR		0.16	20/125	0.8	6/38	39%		ONRKD
CZRHS		0.20	20/100	0.7	6/30	49%		KZVDC
ONHRC		0.25	20/80	0.6	6/24	58%		VSHZO
DKSNV		0.32	20/63	0.5	6/19	67%		HDKCR
ZSOKN		0.40	20/50	0.4	6/15	76%		CSRHN
CKDNR		0.50	20/40	0.3	6/12	80%		SVZDK
SRZKD		0.65	20/32	0.2	6/9.5	84%		NCVOZ
HZOVC		0.80	20/25	0.1	6/7.5	90%		RHSDV
NVDOK		1.00	20/20	0.0	6/6	100%		SNROH
VHCNO		1.20	20/16	-0.1	6/4.8			ODHKR
SVHCZ		1.67	20/12.5	-0.2	6/3.8			ZKCSN
				-0.3				

Der Sehtest ist bei Amlyopiepatienten mit bestmöglicher Korrektur durchzuführen!